
Initiated: 11/73
Reviewed/revised: 8/1/13
Revision: 29

MILWAUKEE
STANDAF
CARDIA

MILWAUKEE COUNTY EMS STANDARD OF CARE CARDIAC ARREST

Approved by: M. Riccardo Colella, DO, MPH, FACEP
Page 1 of 1



NOTES:

- BLS shall be started on all patients in cardiac arrest with the exception of victims with: decapitation; rigor mortis; evidence of tissue decomposition; dependent lividity; presence of a valid Do-Not-Resuscitate or POLST (Physician Orders for Life-Sustaining Treatment); fire victim with full thickness burns to 90% or greater body surface area; hypothermic patients with signs of frozen tissue, rigid airway, ice formation in mouth, or chest noncompliant for CPR.
 - A responding paramedic may cease a BLS initiated resuscitation attempt if:
 - No treatment other than CPR, non-visualized airway insertion, and/or AED application with no shock advised OR patient is in traumatic arrest and ECG shows asystole or PEA at a rate less than 30 OR core temperature is less than 10 °C or 50 °F.
 - If the patient does not meet the above criteria, and a resuscitation attempt is initiated, medical consultation is required to terminate the attempt regardless of the circumstances.
- Resuscitation must be attempted in traumatic cardiac arrests if the patient is in Vfib (defibrillate once and transport) or if the patient has a narrow QRS complex, regardless of the rate.
- The system standard is: CPR will be provided whenever patient is pulseless; compressions at least 100/minute; hands on chest more than 75% of time; minimum compression depth of 2 inches in adults 75% of the time.
- If a fire victim has ROSC, hypotension or altered consciousness, evaluate for possibility of cyanide poisoning and consider administration of hydroxocobalamin (refer to Cyanide Poisoning protocol).
- Please call the Research Line at 805-6493 to report all cardiac arrests, including DOA.